

Note to Readers: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Chronic Exposure to Arsenic and Markers of Cardiometabolic Risk—A Cross-Sectional Study in Chihuahua, Mexico

Michelle A. Mendez, Carmen González-Horta, Blanca Sánchez-Ramírez, Lourdes Ballinas-Casarrubias, Roberto Hernández Cerón, Damián Viniegra Morales, Francisco A. Baeza Terrazas, María C. Ishida, Daniela S. Gutiérrez-Torres, R. Jesse Saunders, Zuzana Drobná, Rebecca C. Fry, John B. Buse, Dana Loomis, Gonzalo G. García-Vargas, Luz M. Del Razo, and Miroslav Stýblo

Table of Contents

Table S1. Urinary As metabolites by quartile of total speciated urinary As: Median (25th-75th percentile).

Table S2. Associations between water and urinary arsenic and continuous indicators of cardiometabolic risk: Coefficient 95% CI.

Table S3. Joint effect of water arsenic combined with arsenic metabolism indicators on cardiometabolic risk: Odds ratios (95% CI).