

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehponline@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Prenatal Exposure to Glycol Ethers and Neurocognitive Abilities in 6-Year-Old Children: The PELAGIE Cohort Study

Rémi Béranger, Ronan Garlantézec, Gaïd Le Maner-Idrissi, Agnès Lacroix, Florence Rouget, Jessica Trowbridge, Charline Warembourg, Christine Monfort, Florent Le Gléau, Marylène Jourdin, Luc Multigner, Sylvaine Cordier, and Cécile Chevrier

Table of Contents

Table S1. Characteristics of the study population compared to the rest of the cohort

Figure S1. Relation between log-transformed GE metabolite concentrations in mothers' prenatal urine samples and the WISC domains of their 6-year-old children

Table S2. Correlation between GE metabolites in mother's prenatal urine samples

Table S3. Associations between glycol ether metabolites and the WISC-IV scores of 6-year-old children in participants with no missing value for covariates (sensitivity analysis)

Table S4. Associations between PhAA concentrations in mothers' prenatal urine samples and the WISC-VCI domain of their 6-year-old children, after exclusion of statistical outliers (sensitivity analysis)

Figure S2. Relation between log-transformed PhAA concentrations in mothers' prenatal urine samples and the WISC-VCI domain of their 6-year-old children, after exclusion of statistical outliers (sensitivity analysis)

Table S5. Associations between concentrations of glycol ether metabolites and visuospatial skills (NEPSY) of 6-year-old children in participants with no missing values for covariates (sensitivity analysis)