

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Particulate Matter and Subclinical Atherosclerosis: Associations between Different Particle Sizes and Sources with Carotid Intima-Media Thickness in the SAPALDIA Study

Inmaculada Aguilera, Julia Dratva, Seraina Caviezel, Luc Burdet, Eric de Groot, Regina E. Ducret-Stich, Marloes Eeftens, Dirk Keidel, Reto Meier, Laura Perez, Thomas Rothe, Emmanuel Schaffner, Arno Schmit-Trucksäss, Ming-Yi Tsai, Christian Schindler, Nino Künzli, and Nicole Probst-Hensch

Table of Contents

List of SAPALDIA investigators

Additional methodological details

Table S1. Characteristics of the SAPALDIA study subjects aged 50 years or older included in the study and those excluded

Figure S1. Scatter plots of log-transformed mean CIMT values by air pollutant

Table S2. Pearson correlation coefficients between air pollution exposure estimates

Table S3. Estimated percent change in CIMT (95% CI) associated with the residuals obtained from: (a) regressing PNC estimates against PM₁₀, PM_{2.5} and LDSA estimates, and (b) regressing LDSA estimates against PM₁₀, PM_{2.5} and PNC estimates

Table S4. Associations between an interdecile range increase in air pollution exposures and CIMT > cohort-specific 75th percentile

Figure S2. Comparative analysis of percent change in CIMT (95% CI) associated

with an interdecile range increase in exposure estimates using three sources of CIMT data from two reading centres

Table S5. Estimated percent change in CIMT (95% CI) associated with a $10 \mu\text{g}/\text{m}^3$ increase in PM_{10} and $\text{PM}_{2.5}$ exposures

Figure S3. Estimated percent change in CIMT (95% CI) associated with a $10 \mu\text{g}/\text{m}^3$ increase in PM_{10} and $\text{PM}_{2.5}$ exposures within subgroups of selected covariates

References