

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

### **Epigenome-Wide Meta-Analysis of Methylation in Children Related to Prenatal NO<sub>2</sub> Air Pollution Exposure**

Olena Gruzieva, Cheng-Jian Xu, Carrie V. Breton, Isabella Annesi-Maesano, Josep M. Antó, Charles Auffray, Stéphane Ballereau, Tom Bellander, Jean Bousquet, Mariona Bustamante, Marie-Aline Charles, Yvonne de Kluizenaar, Herman T. den Dekker, Liesbeth Duijts, Janine F. Felix, Ulrike Gehring, Mònica Guxens, Vincent V.W. Jaddoe, Soesma A. Jankipersadsing, Simon Kebede Merid, Juha Kere, Ashish Kumar, Nathanael Lemonnier, Johanna Lepeule, Wenche Nystad, Christian Magnus Page, Sviatlana Panasevich, Dirkje Postma, Rémy Slama, Jordi Sunyer, Cilla Söderhäll, Jin Yao, Stephanie J. London, Göran Pershagen, Gerard H. Koppelman, and Erik Melén

#### **Table of Contents**

##### **Materials and Methods**

**Table S1.** Characteristics of the included cohorts with available cord blood samples.

**Figure S1.** Plot of the cumulative distribution function for methylation intensities for the three top-ranked CpG sites.

**Figure S2.** Quantile-quantile plot (A) and Manhattan plot (B) for epigenome-wide meta-analysis of the association between NO<sub>2</sub> exposure during pregnancy and cord blood DNA methylation, additionally adjusted for cell type composition.

**Table S2.** Top 25 CpGs from the epigenome-wide meta-analysis of the association between NO<sub>2</sub> exposure during pregnancy and cord blood DNA methylation additionally adjusted for cell composition (n=1,508 newborns from MeDALL, Generation R, CHS and MoBa cohorts).

**Table S3.** Top 25 CpGs from the analysis of the association between NO<sub>2</sub> exposure during pregnancy and cord blood DNA methylation without adjustment for cell composition (left-side part) and with adjustment for cell composition according to Bakulski (right-side) (n=280 newborns from the MeDALL study).

**Table S4.** Look-up in older children for top 25 findings for prenatal NO<sub>2</sub> exposure in relation to methylation in meta-analysis of newborns.

**Table S5.** Look-up in 4-year-old children with paired cord blood and 4-year-old samples for top 25 findings for prenatal NO<sub>2</sub> exposure in relation to methylation in meta-analysis of newborns.

**Table S6.** Nominally significant CpGs within oxidative stress genes extracted from the epigenome-wide meta-analysis of the association between prenatal NO<sub>2</sub> exposure and newborn cord blood DNA methylation (n=1,508 newborns from MeDALL, Generation R, CHS and MoBa cohorts).

**Table S7.** All available CpGs mapped to *CAT* gene from the epigenome-wide meta-analysis of the association between NO<sub>2</sub> exposure during pregnancy and newborn cord blood DNA methylation (N=1,508 newborns from MeDALL, Generation R, CHS and MoBa cohorts).

**Table S8.** All available CpGs mapped to *TPO* gene from the epigenome-wide meta-analysis of the association between NO<sub>2</sub> exposure during pregnancy and newborn cord blood DNA methylation (N=1,508 newborns from MeDALL, Generation R, CHS and MoBa cohorts).

**Table S9.** Look-up in older children for top CpGs in the *CAT* and *TPO* genes for prenatal NO<sub>2</sub> exposure in relation to methylation in meta-analysis of newborns (n=1,508).

**Figure S3.** Regional plots and co-methylation patterns for the genes containing FDR-significant EWAS CpG sites.

## **Acknowledgments**

## **Funding**